

Kites and representations of pseudo MV-algebras

Michał Botur, Tomasz Kowalski

September 10, 2022

Residuated lattices

We work within the general framework of *residuated lattices*, that is, algebras $\mathbf{A} = (A; \wedge, \vee, \cdot, \backslash, /, 1)$ such that

(RL1) $(A; \wedge, \vee)$ is a lattice,

(RL2) $(A; \cdot, 1)$ is a monoid,

(RL3) the equivalences

$$y \leq x \backslash z \quad \Leftrightarrow \quad xy \leq z \quad \Leftrightarrow \quad x \leq z / y$$

hold for all $x, y, z \in A$.

FL-algebras

Expansions of residuated lattices by an additional constant 0 are known as *FL-algebras* and it is there only to make it possible to define *negations*, that is, the operations

$$x^- := 0 / x \quad \text{and} \quad x^\sim := x \setminus 0.$$

FL-algebras

Expansions of residuated lattices by an additional constant 0 are known as *FL-algebras* and it is there only to make it possible to define *negations*, that is, the operations

$$x^- := 0 / x \quad \text{and} \quad x^\sim := x \setminus 0.$$

An FL-algebra **A** is *integral* if 1 is the largest element of A ; it is *0-bounded* if 0 is the smallest element of A . Integral 0-bounded FL-algebras are known as FL_w -algebras (w for *weakening*), so according to our conventions FL_w will stand for the variety of FL_w -algebras.

ℓ -groups and some derived residuated lattices

Definition

A *lattice ordered group* (ℓ -group) is an algebra

$\mathbf{L} = (L; \wedge, \vee, \cdot, ^{-1}, e)$ where $(L; \wedge, \vee)$ is a lattice, $(L; \cdot, ^{-1}, e)$ is a group and

$$x(y \wedge z)w = (xyw) \wedge (xzw),$$

$$x(y \vee z)w = (xyw) \vee (xzw)$$

hold for any $x, y, z, w \in L$.

ℓ -groups and some derived residuated lattices

For our purposes here, it will suffice to recall that any ℓ -group \mathbf{L} is completely determined by the residuation structure of its negative cone $L^- = \{x \in L : x \leq e\}$. Namely, defining the algebra

$$\mathbf{L}^- = (L^-; \wedge, \vee, \cdot, \backslash, /, e)$$

where e , \wedge , \vee and \cdot are inherited from \mathbf{L} , and

$$x / y := (xy^{-1}) \wedge e, \quad y \backslash x := (y^{-1}x) \wedge e$$

ℓ -groups and some derived residuated lattices

For our purposes here, it will suffice to recall that any ℓ -group \mathbf{L} is completely determined by the residuation structure of its negative cone $L^- = \{x \in L : x \leq e\}$. Namely, defining the algebra

$$\mathbf{L}^- = (L^-; \wedge, \vee, \cdot, \backslash, /, e)$$

where e , \wedge , \vee and \cdot are inherited from \mathbf{L} , and

$$x / y := (xy^{-1}) \wedge e, \quad y \backslash x := (y^{-1}x) \wedge e$$

we obtain an integral residuated lattice satisfying the identities

$$xy / y = x = y \backslash yx \quad (\text{Can})$$

$$x / (y \backslash x) = x \vee y = (x / y) \backslash x. \quad (\text{Łuk})$$

The first of these is equivalent over residuated lattices to the usual cancellation laws

$$zx = zy \Rightarrow x = y \quad \text{and} \quad xz = yz \Rightarrow x = y.$$

The second amounts to a non-commutative rendering of the *Łukasiewicz axiom* $(x \rightarrow y) \rightarrow y = (y \rightarrow x) \rightarrow x$.

ℓ -groups and some derived residuated lattices

Members of the variety CanIGMV of *cancellative integral generalised MV-algebras* are (up to isomorphism) precisely the negative cones of ℓ -groups. Namely, there exists functors

$$- : \text{LG} \rightarrow \text{CanIGMV} \text{ and } \ell : \text{CanIGMV} \rightarrow \text{LG}$$

such that $\ell(\mathbf{A})^- = \mathbf{A}$ for any $\mathbf{A} \in \text{CanIGMV}$ and $\ell(\mathbf{L}^-) \cong \mathbf{L}$ for any $\mathbf{L} \in \text{LG}$. These functors establish a categorical equivalence between LG and CanIGMV . In particular, the subvariety lattices of LG and CanIGMV are isomorphic.

Pseudo MV-algebras

Pseudo MV-algebras were originally defined and studied by Georgescu and Iorgulescu, as algebras $(A; \oplus, ^-, \sim, 0, 1)$ satisfying the identities:

$$(A1) \quad x \oplus (y \oplus z) = (x \oplus y) \oplus z,$$

$$(A2) \quad x \oplus 0 = x,$$

$$(A3) \quad x \oplus 1 = 1,$$

$$(A4) \quad (x^- \oplus y^-)^\sim = (x^\sim \oplus y^\sim)^-,$$

$$(A5) \quad (x \oplus y^\sim)^- \oplus x^- = y \oplus (x^- \oplus y)^\sim,$$

$$(A6) \quad x \oplus (y^- \oplus x)^\sim = y \oplus (x^- \oplus y)^\sim$$

$$(A7) \quad x^{-\sim} = x,$$

$$(A8) \quad 0^- = 1.$$

The identities $0 \oplus x = x$ and $1 \oplus x = 1$ follow, as well as $1^- = 0 = 1^\sim$, and $x^{\sim-} = x$.

Pseudo MV-algebras

In any pseudo MV-algebra defined by (A1)–(A8), the lattice operations, multiplication and residuals are defined by

- ▶ $x \vee y := x \oplus (y \odot x^{\sim})$ and $x \wedge y := (x^{-} \oplus y) \odot x$,
- ▶ $x \cdot y := (x^{-} \oplus y^{-})^{\sim}$,
- ▶ $x \setminus y := y \oplus x^{\sim}$ and $y / x := x^{-} \oplus y$.

Pseudo MV-algebras

In any pseudo MV-algebra defined by (A1)–(A8), the lattice operations, multiplication and residuals are defined by

- ▶ $x \vee y := x \oplus (y \odot x^\sim)$ and $x \wedge y := (x^- \oplus y) \odot x$,
- ▶ $x \cdot y := (x^- \oplus y^-)^\sim$,
- ▶ $x \setminus y := y \oplus x^\sim$ and $y / x := x^- \oplus y$.

Then pseudo MV-algebras are (equivalently) bounded IGMV-algebras and the variety ΨMV of *pseudo MV-algebras* is a subvariety of FL_w

Perfect FL_w -algebras

Perfect FL_w -algebras

Definition

An FL_w -algebra \mathbf{A} is perfect if there is a homomorphism $h_{\mathbf{A}}: \mathbf{A} \rightarrow \mathbf{2}$ such that for any $x \in h_{\mathbf{A}}^{-1}(0)$ and any $y \in h_{\mathbf{A}}^{-1}(1)$ the inequality $x \leq y$ holds.

Perfect FL_w -algebras

Definition

An FL_w -algebra \mathbf{A} is perfect if there is a homomorphism $h_{\mathbf{A}}: \mathbf{A} \rightarrow \mathbf{2}$ such that for any $x \in h_{\mathbf{A}}^{-1}(0)$ and any $y \in h_{\mathbf{A}}^{-1}(1)$ the inequality $x \leq y$ holds.

Lemma

Let \mathbf{A} be a perfect FL_w -algebra. Then the homomorphism $f_{\mathbf{A}}: \mathbf{A} \rightarrow \mathbf{2}$ is unique. Hence \mathbf{A} has a unique maximal normal filter.

Perfect FL_w -algebras

For any class \mathcal{K} of FL_w -algebras, we denote the subclass of all its perfect members by \mathcal{K}_{pf} . We say that a variety \mathcal{V} of FL_w -algebras is *perfectly generated* if it is generated by its perfect members, that is, if $\mathcal{V} = V(\mathcal{V}_{pf})$.

Theorem

A subvariety \mathcal{V} of FL_w is perfectly generated if and only if \mathcal{V} is nontrivial and satisfies the following identities:

$$\alpha(x / x^-) \vee \beta(x^- / x) = 1, \quad (1)$$

$$\alpha((x \vee x^-) \cdot (y \vee y^-))^- \leq \alpha((x \vee x^-) \cdot (y \vee y^-)), \quad (2)$$

$$x \wedge x^- \leq y \vee y^- \quad (3)$$

for every $\mathbf{A} \in \mathcal{V}$ and all $\alpha, \beta \in \text{cPol}(\mathbf{A})$.

Perfect FL_w -algebras

Corollary

Any nontrivial subvariety of a perfectly generated variety is also perfectly generated.

Since the trivial variety is not perfectly generated, the variety BA of Boolean algebras is the smallest perfectly generated variety. Indeed, perfectly generated varieties form a lattice ideal in the lattice $\Lambda^+(\text{FL}_w)$, as we will now show.

Theorem

Perfectly generated varieties form an ideal in $\Lambda^+(\text{FL}_w)$.

Kites

Definition

Let \mathbf{L} be an ℓ -group and $\lambda: \mathbf{L} \rightarrow \mathbf{L}$ be an automorphism. We define the algebra

$$\mathcal{K}(\mathbf{L}, \lambda) := (L^- \uplus L^+; \wedge, \vee, \odot, \backslash, /, 0, 1)$$

where $L^- \uplus L^+$ is a disjoint union, $0 := e \in L^+$, $1 := e \in L^-$, and the other operations are given by

$$x \wedge y := \begin{cases} x \wedge y \in L^- & \text{if } x, y \in L^-, \\ x \in L^+ & \text{if } x \in L^+, y \in L^- \\ y \in L^+ & \text{if } x \in L^-, y \in L^+, \\ x \wedge y \in L^+ & \text{if } x, y \in L^+, \end{cases}$$

$$x \vee y := \begin{cases} x \vee y \in L^- & \text{if } x, y \in L^-, \\ y \in L^- & \text{if } x \in L^+, y \in L^-, \\ x \in L^- & \text{if } x \in L^-, y \in L^+, \\ x \vee y \in L^+ & \text{if } x, y \in L^+, \end{cases}$$

$$x \odot y := \begin{cases} x \cdot y \in L^- & \text{if } x, y \in L^-, \\ \lambda(x) \cdot y \vee e \in L^+ & \text{if } x \in L^-, y \in L^+, \\ x \cdot y \vee e \in L^+ & \text{if } x \in L^+, y \in L^-, \\ e \in L^+ & \text{if } x, y \in L^+, \end{cases}$$

$$\begin{aligned}
x \setminus y &:= \begin{cases} x^{-1} \cdot y \wedge e \in L^{-} & \text{if } x, y \in L^{-}, \\ e \in L^{-} & \text{if } x \in L^{+}, y \in L^{-} \\ \lambda(x)^{-1} \cdot y \vee e \in L^{+} & \text{if } x \in L^{-}, y \in L^{+}, \\ x^{-1} \cdot y \wedge e \in L^{-} & \text{if } x, y \in L^{+}, \end{cases} \\
y / x &:= \begin{cases} y \cdot x^{-1} \wedge e \in L^{-} & \text{if } x, y \in L^{-}, \\ e \in L^{-} & \text{if } x \in L^{+}, y \in L^{-} \\ y \cdot x^{-1} \vee e \in L^{+} & \text{if } x \in L^{-}, y \in L^{+}, \\ \lambda^{-1}(y \cdot x^{-1}) \wedge e \in L^{-} & \text{if } x, y \in L^{+}, \end{cases}
\end{aligned}$$

Kites

Now, in any perfect pseudo MV-algebra \mathbf{A} the normal filter $F_{\mathbf{A}}$ is the universe of a cancellative IGMV-algebra $\mathbf{F}_{\mathbf{A}}$. Since pseudo MV-algebras satisfy the identities

$$(x \wedge y)^{\sim\sim} = x^{\sim\sim} \wedge y^{\sim\sim}$$

$$(x \vee y)^{\sim\sim} = x^{\sim\sim} \vee y^{\sim\sim}$$

$$(x \cdot y)^{\sim\sim} = x^{\sim\sim} \cdot y^{\sim\sim}$$

$$x^{-\sim\sim} = x^{\sim\sim-}$$

the map $-\sim\sim$ is an automorphism of $\mathbf{F}_{\mathbf{A}}$. Applying the functor ℓ we lift $-\sim\sim$ to an automorphism

$$\ell^{\sim} : \ell(\mathbf{F}_{\mathbf{A}}) \rightarrow \ell(\mathbf{F}_{\mathbf{A}})$$

defined, obviously, as $\ell^{\sim}(-) := \ell(-\sim\sim)$.

Theorem

Let \mathbf{A} be a perfect pseudo MV-algebra. Then $\mathbf{A} \cong \mathcal{K}(\ell(\mathbf{F}_{\mathbf{A}}), \ell^{\sim})$.

Kites

Definition

We define \mathbf{LGA} to be the category of ℓ -groups with a distinguished automorphism. The objects are algebras (\mathbf{L}, λ) where \mathbf{L} is an ℓ -group and λ is an automorphism of \mathbf{L} . The morphisms are ℓ -group homomorphisms commuting with the distinguished automorphism.

Theorem

The categories $\mathbf{pf}\Psi\mathbf{MV}$ of perfect pseudo MV-algebras, and \mathbf{LGA} of ℓ -groups with a distinguished automorphism, are equivalent.

Corollary (Di Nola, Lettieri)

Let \mathbf{AbLG} be the category of Abelian ℓ -groups with homomorphisms and let \mathbf{pfMV} be the category of perfect MV-algebras with homomorphisms. Then \mathbf{AbLG} and \mathbf{pfMV} are equivalent.

Definition

A monounary algebra $\mathbf{B} = (B; \beta)$ where β is a bijection on B will be called a B -cycle. Homomorphisms of B -cycles are maps $f: \mathbf{B} \rightarrow \mathbf{C}$ satisfying $f \circ \lambda^{\mathbf{B}} = \lambda^{\mathbf{C}} \circ f$. Objects of the category \mathbf{BC} are B -cycles and arrows are homomorphisms.

Remark

B -cycles are not a variety, but as we will often need β^{-1} , we could have equivalently defined B -cycles as a variety of bi-unary algebras (B, β, δ) satisfying $\beta(\delta(x)) = x = \delta(\beta(x))$, and write β^{-1} for δ .

Definition

Let $\mathbf{B} = (B; \beta)$ be a B -cycle and \mathbf{L} and ℓ -group. A kite over \mathbf{B} and \mathbf{L} is the algebra

$$\mathcal{K}_{\mathbf{B}}(\mathbf{L}) := \mathcal{K}(\mathbf{L}^B, \lambda)$$

where $\lambda: \mathbf{L}^B \rightarrow \mathbf{L}^B$ is the automorphism given by $\lambda(x(i)) = x(\beta(i))$ for any $i \in B$.

Let \mathbf{L} be an ℓ -group, and let $f: \mathbf{B} \rightarrow \mathbf{C}$ be a homomorphism of B-cycles. Then f naturally lifts to a homomorphism

$$\mathcal{K}_f(\mathbf{L}): \mathcal{K}_{\mathbf{C}}(\mathbf{L}) \rightarrow \mathcal{K}_{\mathbf{B}}(\mathbf{L})$$

defined by $(\mathcal{K}_f(\mathbf{L}))(x) = x \circ f$. Moreover

$$\mathcal{K}_-(\mathbf{L}): \mathbf{BC} \rightarrow \mathbf{pf}\Psi\mathbf{MV}$$

is a contravariant functor.

Definition

For a perfect MV-algebra \mathbf{A} and an ℓ -group \mathbf{L} we define a B-cycle $[\mathbf{A}, \mathbf{L}] := ([A, L]; \lambda)$, where

$$[A, L] := \{\alpha \in (L^-)^{F_{\mathbf{A}}} : \alpha \text{ is a homomorphism in } \text{CanIGMV}\}$$

given by $\lambda(\alpha(x)) := \alpha(x^{\sim\sim})$ for any $\alpha \in [A, L]$.

For any perfect pseudo MV-algebra \mathbf{A} there exists a homomorphism

$$\eta_{\mathbf{A}}: \mathbf{A} \rightarrow \mathcal{K}_{[\mathbf{A}, \mathbf{L}]}(\mathbf{L}).$$

(defined by

$$(\eta_{\mathbf{A}}(x))(\alpha) = \begin{cases} \alpha(x) & \text{if } x \in F_{\mathbf{A}}, \\ \alpha(x^{\sim})^{-1} & \text{if } x \in C_{\mathbf{A}}. \end{cases}$$

for any $x \in |\mathbf{A}|$ and any $\alpha \in [A, L]$)

For any perfect pseudo MV-algebra \mathbf{A} there exists a homomorphism

$$\eta_{\mathbf{A}}: \mathbf{A} \rightarrow \mathcal{K}_{[\mathbf{A}, \mathbf{L}]}(\mathbf{L}).$$

(defined by

$$(\eta_{\mathbf{A}}(x))(\alpha) = \begin{cases} \alpha(x) & \text{if } x \in F_{\mathbf{A}}, \\ \alpha(x^{\sim})^{-1} & \text{if } x \in C_{\mathbf{A}}. \end{cases}$$

for any $x \in |\mathbf{A}|$ and any $\alpha \in [A, L]$)

Theorem

Let \mathbf{A} be a perfect pseudo MV-algebra, \mathbf{B} be a B-cycle and \mathbf{L} be an ℓ -group. Then for any homomorphism $f: \mathbf{A} \rightarrow \mathcal{K}_{\mathbf{B}}(\mathbf{L})$ there exists a unique homomorphism of B-cycles $g: \mathbf{B} \rightarrow [\mathbf{A}, \mathbf{L}]$ such that the following diagram commutes:

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{\eta_{\mathbf{A}}} & \mathcal{K}_{[\mathbf{A}, \mathbf{L}]}(\mathbf{L}) \\ & \searrow f & \downarrow \kappa_g(\mathbf{L}) \\ & & \mathcal{K}_{\mathbf{B}}(\mathbf{L}). \end{array}$$

Varieties generated by Dvurečenskij/Kowalski's kites

Throughout the section \mathbb{D} will stand for the lattice $(\mathbb{N}; |)$ of natural numbers under the divisibility ordering.

Varieties generated by Dvurečenskij/Kowalski's kites

Throughout the section \mathbb{D} will stand for the lattice $(\mathbb{N}; |)$ of natural numbers under the divisibility ordering.

For any bijection λ on a nonempty set B , we define the *dimension* of λ as follows:

$$\dim(\lambda) := \min^{\mathbb{D}} \{n \in \mathbb{N} : \lambda^n = id_B\}.$$

For a B-cycle $\mathbf{B} = (B; \lambda)$, we put $\dim(\mathbf{B}) := \dim(\lambda)$ and call it the dimension of \mathbf{B} .

Varieties generated by Dvurečenskij/Kowalski's kites

Throughout the section \mathbb{D} will stand for the lattice $(\mathbb{N}; |)$ of natural numbers under the divisibility ordering.

For any bijection λ on a nonempty set B , we define the *dimension* of λ as follows:

$$\dim(\lambda) := \min^{\mathbb{D}} \{n \in \mathbb{N} : \lambda^n = id_B\}.$$

For a B-cycle $\mathbf{B} = (B; \lambda)$, we put $\dim(\mathbf{B}) := \dim(\lambda)$ and call it the dimension of \mathbf{B} .

Lemma

We have $\Lambda^+(\mathbf{BC}) \cong \mathbb{D}$ and $\Lambda(\mathbf{BC}) \cong \mathbf{1} \oplus \mathbb{D}$, that is, the ordinal sum of the trivial lattice $\mathbf{1}$ and \mathbb{D} .

Varieties generated by Dvurečenskij/Kowalski's kites

Throughout the section \mathbb{D} will stand for the lattice $(\mathbb{N}; |)$ of natural numbers under the divisibility ordering.

For any bijection λ on a nonempty set B , we define the *dimension* of λ as follows:

$$\dim(\lambda) := \min^{\mathbb{D}} \{n \in \mathbb{N} : \lambda^n = id_B\}.$$

For a B-cycle $\mathbf{B} = (B; \lambda)$, we put $\dim(\mathbf{B}) := \dim(\lambda)$ and call it the dimension of \mathbf{B} .

Lemma

We have $\Lambda^+(\mathbf{BC}) \cong \mathbb{D}$ and $\Lambda(\mathbf{BC}) \cong \mathbf{1} \oplus \mathbb{D}$, that is, the ordinal sum of the trivial lattice $\mathbf{1}$ and \mathbb{D} .

We will write C_n for the variety defined by $\lambda^n(x) = x$, so in particular $\mathbf{BC} = C_0$.

Varieties generated by kites

For any pseudo MV-algebra \mathbf{A} , the operation $-\approx$ is a bijection on A , so for any \mathbf{A} we define the dimension of \mathbf{A} to be $\dim(-\approx)$. This is essential for the rest of the section, so we state it formally.

Definition

Let $\mathbf{A} \in \mathbf{PMV}$ and $\mathcal{V} \in \mathbf{\Psi MV}$. Then

1. $\dim(\mathbf{A}) := \dim(-\approx)$,
2. $\dim(\mathcal{V}) := \min^{\mathbb{D}}\{\dim(\mathbf{A}) \mid n : \text{for all } \mathbf{A} \in \mathcal{V}\}$,
3. $\mathbf{P\Psi MV}_n := \mathbf{P\Psi MV} \cap \text{Mod}\{\lambda^n(x) = x\}$, for any $n \in \mathbb{D}$.

Varieties generated by kites

Definition

We define two pairs of maps

$$\begin{aligned}\psi &: \Lambda(\mathbf{P}\Psi\mathbf{MV}) \rightarrow \Lambda(\mathbf{CanI}\mathbf{GMV}), \text{ where } \psi(\mathcal{V}) = V\{\mathbf{F}_{\mathbf{A}} : \mathbf{A} \in \mathcal{V}_{pf}\}, \\ \Psi &: \Lambda(\mathbf{P}\Psi\mathbf{MV}) \rightarrow \Lambda(\mathbf{CanI}\mathbf{GMV}) \times \mathbb{D}, \text{ where } \Psi(\mathcal{V}) = (\psi(\mathcal{V}), \dim(\mathcal{V})),\end{aligned}$$

for any $\mathcal{V} \in \Lambda(\mathbf{P}\Psi\mathbf{MV})$ and

$$\begin{aligned}\delta &: \Lambda(\mathbf{CanI}\mathbf{GMV}) \rightarrow \Lambda(\mathbf{P}\Psi\mathbf{MV}), \text{ where } \delta(\mathcal{V}) = V\{\mathbf{A} \in \mathbf{pf}\Psi\mathbf{MV} : \mathbf{F}_{\mathbf{A}} \in \mathcal{V}\} \\ \Delta &: \Lambda(\mathbf{CanI}\mathbf{GMV}) \times \mathbb{D} \rightarrow \Lambda(\mathbf{P}\Psi\mathbf{MV}), \text{ where } \Delta(\mathcal{V}, n) = \delta(\mathcal{V}) \cap \mathbf{P}\Psi\mathbf{MV}_n,\end{aligned}$$

for any $\mathcal{V} \in \Lambda(\mathbf{CanI}\mathbf{GMV})$ and $n \in \mathbb{D}$.

Varieties generated by kites

Lemma

Let $\mathcal{V} \in \Lambda(\text{CanlGMV})$, let E be an equational base for \mathcal{V} , and let $\mathbf{A} \in \text{P}\Psi\text{MV}$. The following are equivalent.

1. $\mathbf{A} \in \delta(\mathcal{V})$,
2. $\mathbf{A} \models t(x_1 \vee x_1^-, \dots, x_k \vee x_k^-) = 1$ for all terms t in the language of residuated lattices, such that $\mathcal{V} \models t(x_1, \dots, x_k) = 1$.
3. $\mathbf{A} \models t_\varepsilon(x_1 \vee x_1^-, \dots, x_k \vee x_k^-) = 1$ for all equations $\varepsilon(x_1, \dots, x_k) \in E$.

Varieties generated by kites

Theorem

For any $\mathcal{V} \in \Lambda^+(\text{CanlGMV})$ and any $n \in \mathbb{D}$, we have

$$(\mathcal{V}, n) = \Psi\Delta(\mathcal{V}, n).$$

Lemma

For any $\mathcal{V} \in \Lambda(\text{P}\Psi\text{MV})$ we have

$$\mathcal{V} \subseteq \Delta\Psi(\mathcal{V}).$$

Varieties generated by kites

Theorem

For any $\mathcal{V} \in \Lambda^+(\text{CanIGMV})$ and any $n \in \mathbb{D}$, we have

$$(\mathcal{V}, n) = \Psi\Delta(\mathcal{V}, n).$$

Lemma

For any $\mathcal{V} \in \Lambda(\text{P}\Psi\text{MV})$ we have

$$\mathcal{V} \subseteq \Delta\Psi(\mathcal{V}).$$

Theorem

Let $\mathcal{V} \in \Lambda(\text{P}\Psi\text{MV})$. The following are equivalent.

- 1. \mathcal{V} is generated by kites.*
- 2. $\mathcal{V} = \Delta\Psi(\mathcal{V})$.*
- 3. $\mathcal{V} = \Delta(\mathcal{W}, n)$ for some $\mathcal{W} \in \Lambda(\text{CanIGMV})$ and some $n \in \mathbb{D}$.*

Theorem

Varieties generated by kites form a complete sublattice of $\Lambda(\text{P}\Psi\text{MV})$ with $\text{P}\Psi\text{MV}$ being its largest, and BA its smallest element.

Theorem

Let \mathbb{K} be the lattice of subvarieties of $\text{P}\Psi\text{MV}$ generated by kites.

$$\mathbb{K} \cong \mathbf{1} \oplus (\Lambda^+(\text{CanIGMV}) \times \mathbb{D}) \cong \mathbf{1} \oplus (\Lambda^+(\text{LG}) \times \mathbb{D})$$

where $\mathbf{1}$ is the trivial lattice and \oplus is the operation of ordinal sum.

Thank you for your attention!